



31st Austrian Mathematical Olympiad

Federal Competition for Advanced Students

Day 1, June 7, 2000

1. The sequence $\langle a_n \rangle$ with $a_0 = 4$ and $a_1 = 1$ satisfies the recursion $a_{n+1} = a_n + 6a_{n-1}$ for $n \geq 1$ and defines the sequence

$$b_n = \sum_{k=0}^n \binom{n}{k} a_k.$$

Determine the coefficients α and β such that b_n satisfies the recursion

$$b_{n+1} = \alpha b_n + \beta b_{n-1}.$$

Furthermore, determine an explicit term for b_n .

2. The trapezoid $ABCD$ ($ABCD$ labeled counterclockwise, $AB \parallel CD$) is inscribed into a circle k . On the arc \widehat{AB} two points P and Q ($P \neq Q$) are chosen (with $APQB$ labeled in counterclockwise order).

Let X be the intersection of the lines CP and AQ and Y the of intersection of the lines BP and DQ .

Show that P , Q , X and Y lie on a circle.

3. Determine all real solutions of the equation

$$||| ||| |x^2 - x - 1| - 3| - 5| - 7| - 9| - 11| - 13| = x^2 - 2x - 48.$$



31st Austrian Mathematical Olympiad
Federal Competition for Advanced Students
Day 2, June 8, 2000

4. In the acute-angled, non-isosceles triangle $\triangle ABC$ with angle $\gamma = 60^\circ$ let U be the circumcenter, H the orthocenter and D the intersection of the lines AH and BC (that is, the orthogonal projection of A onto BC)

Show that the Euler line HU is the bisector of the angle $\angle BHD$.

5. Determine all pairs of two integers (m, n) such that

$$|(m^2 + 2000m + 999999) - (3n^3 + 9n^2 + 27n)| = 1$$

holds.

6. Determine all functions f mapping from the set of real numbers to the set of real numbers, such that for all real numbers x, y, z the relation $f(x + f(y + z)) + f(f(x + y) + z) = 2y$ holds.