



37th Austrian Mathematical Olympiad
Federal Competition for Advanced Students
Part 1, May 21, 2006

1. Let n be a non-negative integer whose decimal expansion ends with exactly k zeros but that is larger than 10^k .

For a certain n , only $k = k(n) \geq 2$ is known. How many possibilities (as a function of $k = k(n)$) are there at least to represent n as the difference of the squares of two non-negative integers?

2. Show that the sequence $\left\langle \frac{(n+1)^n n^{2-n}}{7n^2+1} \right\rangle_{n=0,1,2,\dots}$ is strictly increasing.
3. In a triangle ABC let D and E be points where the incircle touches the sides BC and AC . Show that if $\overline{AD} = \overline{BE}$, then the triangle is isosceles.
4. We are given the function $f(x) = \lfloor x^2 \rfloor + \{x\}$ that is defined for all positive real numbers. (Here for a real number u , $k = \lfloor u \rfloor$ is the largest integer smaller or equal u and $\{u\} = u - \lfloor u \rfloor$.) Show that there exists an infinite arithmetic sequence of distinct positive rational numbers that all have 3 as denominator when fully reduced and are not in the image of the function f .



37th Austrian Mathematical Olympiad

Federal Competition for Advanced Students

Part 2, Day 1, May 31, 2006

1. Let N be a positive integer.

Determine the number of positive integers $n \leq N$ that have an integer multiple whose decimal expansion only contains the digits 2 and 6 (not necessarily equally often).

2. Show that for all triple (a, b, c) of positive real numbers the inequality

$$3(a + b + c) \geq 8\sqrt[3]{abc} + \sqrt[3]{\frac{a^3 + b^3 + c^3}{3}}$$

holds.

When does equality hold?


3. We are given a triangle ABC .

On the extension of AB beyond B we construct a point R with $\overline{BR} = \overline{BC}$ and on the extension of AC beyond C we construct a point S with $\overline{CS} = \overline{CB}$. Let A' be the intersection of the diagonals in the quadrilateral $BRSC$.

Analogously a point T with $\overline{CT} = \overline{CA}$ is constructed on the extension of BC beyond C , and a point U with $\overline{AU} = \overline{AC}$ on the extension of BA beyond A . Let B' be the intersection of the diagonals in the quadrilateral $CTUA$.

Furthermore a point V with $\overline{AV} = \overline{AB}$ is constructed on the extension of CA beyond A , and a point W with $\overline{BW} = \overline{BA}$ on the extension of CB beyond B . Let C' be the intersection of the diagonals in the quadrilateral $AVWB$.

Show that the area $F(AC'BA'CB')$ of the hexagon equals the sum of the areas $F(ABC)$ and $F(A'B'C')$ of the two triangles whose sides are the diagonals of the hexagon.



37th Austrian Mathematical Olympiad
Federal Competition for Advanced Students
Part 2, Day 2, June 1, 2006

4. Determine all rational numbers x such that $1 + 105 \cdot 2^x$ is the square of a rational number.
5. Find all monotonic functions $f: \mathbb{R} \rightarrow \mathbb{R}$ with $f(-f(x)) = f(f(x)) = f(x)^2$.
($f(x)$ is called monotonic if either $a < b \Rightarrow f(a) \leq f(b)$ or $a < b \Rightarrow f(a) \geq f(b)$.)
6. For integers $A \neq 0$ solve the following system of equations over $\mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}$.

$$\begin{aligned}x + y^2 + z^3 &= A \\ \frac{1}{x} + \frac{1}{y^2} + \frac{1}{z^3} &= \frac{1}{A} \\ x \cdot y^2 \cdot z^3 &= A^2\end{aligned}$$