



39th Austrian Mathematical Olympiad

Federal Competition for Advanced Students

Part 1, May 22, 2008

1. What is the smallest positive residue of

$$1 \cdot \binom{2008}{0} + 2 \cdot \binom{2008}{1} + 3 \cdot \binom{2008}{2} + \dots + 2009 \cdot \binom{2008}{2008}$$

when divided by 2008?

2. Let a be a positive real number and $n > 4$ an integer.

Determine all n -tuples (x_1, x_2, \dots, x_n) of positive real numbers that satisfy the following system of equations.

$$\begin{aligned} x_1 x_2 (3a - 2x_3) &= a^3 \\ x_2 x_3 (3a - 2x_4) &= a^3 \\ &\vdots \\ x_{n-2} x_{n-1} (3a - 2x_n) &= a^3 \\ x_{n-1} x_n (3a - 2x_1) &= a^3 \\ x_n x_1 (3a - 2x_2) &= a^3 \end{aligned}$$

3. Let $p > 1$ be a natural number.

We consider the set \mathbf{F}_p of all non-constant sequences of non-negative integers that satisfy the recursion $a_{n+1} = (p+1)a_n - pa_{n-1}$ (for all $n > 0$).

Show that there exists a sequence $\langle a_n \rangle$ in \mathbf{F}_p with the property that for every other sequence $\langle b_n \rangle$ in \mathbf{F}_p , the inequality $a_n \leq b_n$ holds for all n .

4. In a triangle ABC let E be the midpoint of the sides AC and F the midpoint of the side BC . Furthermore let G be the foot of the altitude through C on the side AB (or its extension).

Show that the triangle EFG is isosceles if and only if ABC is isosceles.



39th Austrian Mathematical Olympiad
 Federal Competition for Advanced Students
Part 2, Day 1, June 5, 2008

1. Show that for positive real numbers a, b, c with $a + b + c = 1$ the inequality

$$\sqrt{a^{1-a}b^{1-b}c^{1-c}} \leq \frac{1}{3}$$

holds.

2. (a) Does there exist a polynomial $P(x)$ with integer coefficients such that for each positive integer d that divides 2008, one has $P(d) = \frac{2008}{d}$?
- (b) For which positive integers n do exist polynomials $P(x)$ with integer coefficients such that for each positive integers d that divides n , one has $P(d) = \frac{n}{d}$?
3. The line g is given, and on it the four points P, Q, R, S (in this order from left to right).


Construct all squares $ABCD$ with the following properties:

P lies on the line through A and D .

Q lies on the line through B and C .

R lies on the line through A and B .

S lies on the line through C and D .



39th Austrian Mathematical Olympiad

Federal Competition for Advanced Students

Part 2, Day 2, June 6, 2008

4. Determine all functions f mapping the set of positive integers to the set of non-negative integers satisfying the following conditions:

(a) $f(n \cdot m) = f(m) + f(n)$;

(b) $f(2008) = 0$;

(c) For all $n \equiv 39 \pmod{2008}$ one has $f(n) = 0$.

5. Which positive integers are missing in the sequence $\langle a_n \rangle$?

$$\langle a_n = n + \lfloor \sqrt{n} \rfloor + \lfloor \sqrt[3]{n} \rfloor \rangle_{n \geq 1}.$$

6. We are given a square $ABCD$. Let P be different from the vertices of the square and from its center M .

For a point P for which the line PD intersects the line AC , let E be this intersection.

For a point P for which the line PC intersects the line DB , let F be this intersection.

All those points P for which E and F exist are called acceptable points.

Determine the set of acceptable points for which the line EF is parallel to AD .