



37th Austrian Mathematical Olympiad

Regional Competition for Advanced Students

April 27, 2006

1. Let $0 < x < y$ be real numbers, and let

$$H = \frac{2xy}{x+y}, \quad G = \sqrt{xy}, \quad A = \frac{x+y}{2}, \quad Q = \sqrt{\frac{x^2+y^2}{2}}$$

be the harmonic, geometric, arithmetic and quadratic means of x and y . It is known that $H < G < A < Q$ holds.

Sort the intervals $[H, G]$, $[G, A]$ and $[A, Q]$ ascendingly by their lengths.

2. Let $n > 1$ be a natural number and a a real number.

Determine all real solutions (x_1, x_2, \dots, x_n) of the following system of equations:

$$\begin{aligned} x_1 + ax_2 &= 0 \\ x_2 + a^2x_3 &= 0 \\ &\vdots \\ x_k + a^kx_{k+1} &= 0 \\ &\vdots \\ x_{n-1} + a^{n-1}x_n &= 0 \\ a^n x_1 + x_n &= 0 \end{aligned}$$

3. In a non-isosceles triangle ABC let w be the bisector of the exterior angle in C (exterior bisector of γ). The intersection of w with the extension of AB is D . Let now k_A be the circumcircle of the triangle ADC and analogously k_B the circumcircle of the triangle BDC . Let t_A be the tangent to k_A in A and analogously t_B the tangent to k_B in B . Let P be the intersection of these two tangents.

Let now the points A and B be given.

Determine the set of the points $P = P(C)$ for all points C such that ABC is a non-isosceles, acute-angled triangle.

4. Let $\langle h_n \rangle_{n \in \mathbb{N}}$ be a harmonic sequence of positive rational numbers (i.e., each h_n is the harmonic mean of its neighbors h_{n-1} and h_{n+1} , that is, $h_n = \frac{2h_{n-1}h_{n+1}}{h_{n-1}+h_{n+1}}$).

Show that if the sequence contains an element h_j that is the square of a rational number, then it contains infinitely many elements h_k that are squares of rational numbers.